数学与系统科学研究院 计算数学所学术报告

<u>报告人</u>: Prof. Jianliang Qian

(Department of Mathematics, Michigan State University, USA)

<u>报告题目</u>:

Hadamard-Babich method for point-source Maxwell's equations in inhomogeneous media

<u>邀请人:</u> 陈志明 院士

<u>报告时间</u>: 2018 年 7 月 30 日(周一) 上午 10:00-11:00

<u>报告地点</u>:数学院南楼九层

902 教室

Abstract:

Starting from Hadamard's method, we extend Babich's ansatz to the frequency-domain point-source (FDPS) Maxwell's equations in an inhomogeneous medium in the high-frequency regime. First, we develop a novel asymptotic series, dubbed Hadamard's ansatz, to form the fundamental solution of the Cauchy problem for the time-domain point-source (TDPS) Maxwell's equations in the region Governing equations for the unknowns in close to the source. Hadamard's ansatz are then derived. In order to derive the initial data for the unknowns in the ansatz, we further propose a condition for matching Hadamard's ansatz with the homogeneous-medium fundamental solution at the source. Directly taking the Fourier transform of Hadamard's ansatz in time, we obtain a new ansatz, dubbed the Hadamard-Babich ansatz, for the FDPS Maxwell's equations. Next, we elucidate the relation between the Hadamard-Babich ansatz and a recently proposed Babich-like ansatz for solving the same FDPS Maxwell's equations. Finally, incorporating the first two terms of the Hadamard-Babich ansatz into a planar-based Huygens sweeping algorithm, we solve the FDPS Maxwell's equations at high frequencies in the region where caustics Numerical experiments demonstrate the accuracy of our occur. method.

欢迎大家参加!