

数学与系统科学研究院

计算数学所学术报告

报告人: **Dr. Karolina Kropielnicka**

(*University of Gdańsk and Polish Academy of Sciences, Poland*)

报告题目:

**Exponential splitting, Lie algebra,
and commutator-free Magnus based
methods for the linear Schrödinger
equation**

邀请人: 唐贻发 研究员

报告时间: **2017年6月6日 (周二)**

下午 16:00-17:00

报告地点: 数学院南楼七层

714 教室

Abstract:

The discretization of a linear Schrödinger equation is difficult due to the presence of a small parameter which induces high oscillations. A standard approach consists of a spectral semidiscretization, followed by an exponential splitting. This, however, is sub-optimal, because the exceedingly high precision in space discretization is married by low order of the time solver. In this talk we sketch an alternative approach.

Our analysis commences not with semi-discretisation, but with the investigation of the Lie algebra generated by differentiation and by multiplication with the interaction potential: it turns out that this algebra possesses a structure which renders it amenable to a very effective form of asymptotic splitting: exponential splitting where consecutive terms are scaled by increasing powers of the small parameter. The semi-discretisation is deferred to the very end of computations.

We will focus on the method for the time dependant linear Schrödinger equation with potential non-depending on time, however we will also discuss the difficulties that appear with time dependant potential and will briefly propose the remedy to that stage of an affair, a commutator-free Magnus based method.

Based on joint work with, Philipp Bader (La Trobe University, Australia), Iserles Arieh (University of Cambridge, UK) and Pranav Singh (University of Oxford, UK)

欢迎大家参加！